



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A23L 1/10, 1/168, 1/308, A61K 35/78	A1	(11) International Publication Number: WO 99/08546 (43) International Publication Date: 25 February 1999 (25.02.99)
(21) International Application Number: PCT/US98/17053 (22) International Filing Date: 18 August 1998 (18.08.98) (30) Priority Data: 60/056,562 19 August 1997 (19.08.97) US (71) Applicant (for all designated States except US): BARKLEY SEED INC. [US/US]; P.O. Box 5540, Yuma, AZ 85364 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): FOX, Gregory, J. [US/US]; 717 14th Street South, Fargo, ND 58103 (US). (74) Agents: JONDLE, Robert, J. et al.; Rothwell, Figg, Ernst & Kurz, Suite 701 East, 555 13th Street, N.W., Columbia Square, Washington, DC 20004 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: HIGH VISCOSITY CEREAL AND FOOD INGREDIENT FROM VISCOUS BARLEY GRAIN (57) Abstract <p>Disclosed are highly digestible barley flakes and kibbles which have elevated levels of long chained beta glucan fibers (1 mm to 6 mm long, 2 microns to 10 microns wide) providing these cereal foods and food ingredients with clinically active hypocholesterolaemic, immunostimulative, and atumorigenic properties; also disclosed are the methods to produce these flakes and kibbles.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

High Viscosity Cereal and Food Ingredient from Viscous Barley Grain

Field of Invention

The present invention relates to a clinically active
5 hypocholesterolaemic, immunostimulative, atumorigenic food and/or food
ingredients derived from waxy hulless and waxy hulless short awn barley
varieties, that have elevated levels of long chained beta glucan fiber, which
is the active clinical ingredient in this food product.

It is well know in the cereal and nutritional arts that beta glucan
10 (beta 1-4/beta 1-3 glucosyl pyranose polymer) reduces serum cholesterol,
and regulates serum blood sugar in humans. Beta glucan occurs in
quantity in only two cereals as soluble cell wall fiber, i.e. oats and barley.
Two forms of barley have especially high levels of beta glucan, i.e. waxy
hulless and waxy hulless short awn. The clinical activity of these beta
15 glucans is related to the viscosity that is created in the intestines, and that
this viscosity is created by hydrated beta glucan fibers that ranged from 1
to 6 mm in length and about 2 to 10 microns in diameter. The viscosity is
directly related to the quantity and length of these fibers. These sticky
fibers absorb cholesterol, fat, and sugars and flush them out of the body,
20 this flushing action is the basis of the clinical functionality.

These same beta glucans, which are common as cell wall fiber in
microbes have immunostimulative, and anti-tumorigenic properties. More
recently it has become apparent that the cereal beta glucans have the same

healing properties as the microbial forms. Additional new information strongly suggests that these same beta glucans have anti-inflammatory properties that have a healing effect on bone and joint disorders, and a flushing action of the blood, lymph, and interstitial fluids that results in anti cancer effects. The pharmacological functionality is related to the same properties as the nutritional functionality, chain length of beta glucan fibers. As beta glucan chains pass through the intestinal tract they are broken up in length and absorbed by intestinal lymph nodes, then passing to the other organ systems. As molecular length is reduced, functionality is reduced. Thus, the long chained beta glucan fibers associated with the waxy hulless and waxy hulless short awned barleys would be especially effective in this role as a nutritional pharmacological agent.

In the last 3 years the grain growing regions of the United States have been plagued by a fungal disease called scab, that grows on the surface of the seed, this will effect all seed to some extent, but the seed which are severely effected will be bleached and shrunken, and in the industry are called tombstones because of their appearance. This fungus produces toxins that are collectively called vomitoxins, because in quantities around 10 ppm they will cause pigs to become sick with one of the symptoms, vomiting. The disease is not considered to be toxic to humans, but there is some controversy about this point, because if nothing else people allergic to vomitoxin can die from anaphylactic shock in response to these vomitoxins. At the very least they can make people sick, and certainly are not healthful compounds.

The healthful beta glucans of barley and especially waxy hulless and waxy hulless short awn barley (High Vee™ Barley) are concentrated in the endosperm of the grain, not the bran. This makes barley unique because in oats the beta glucans are concentrated in the bran. While the general

public regards bran as healthful, this is in fact some what of a misnomer in real world practice. In fact diseases like scab grow on the surface of the grain. Also, the bran fraction contains high concentrations of phytic acid, a compound that ties up essential minerals and causes mineral deficiencies in live stock and has the potential to do the same in humans. Also, the harsh lignified fiber that occurs in the bran fraction, if taken in large quantities, can have a detrimental tearing effect on the intestinal cell walls. Thus, the benefit of the fiber associated with the bran fraction is mixed at best. In the present day climate of bran acceptance, as a health benefit, it is forgotten that the development of refined milling procedures in Northern Europe to remove bran from flour was a health measure to remove toxic fungi adhering to the moldy bran, that was inevitable in the cool moist environment of Northern Europe. In waxy hulless barley, and waxy hulless short awn barley there exists a highly effective form of fiber (long chained beta glucan) that is not associated with the potentially moldy bran. In this respect waxy hulless and waxy hulless short awn barleys are truly unique.

If one looks back to antiquity, to the beginnings of human civilization, barley grain was the base component of the first breads. Barley bread was considered the bread of life. As time went on the breads wheats with strong glutinous proteins, which produced the leaven loaves became the preferred bread of the affluent. However, even as these fluffy wheat breads were being introduced, their lack of healthfulness as compared to the old heavy barley breads was noted by numerous sources. These observations continued into the 19th century AD, when high gluten strength wheat breads finally replaced the heavy barley, corn, oat, rye type breads as the bread of the people, not just the affluent. Always it was noted that the raised wheat breads while fun to eat somehow seemed less healthful than the old breads, especially those with a barley base. It

should be noted that the old heavy breads were quite often a combination of grain (multi grain breads). Indeed the Roman Army that conquered much of the known world marched on multi grain bread. However, of the grains in their bread, barley was the one, that in particular was thought to
5 give strength and vigor to the soldiers. Wheat was a luxury, barley was a functional necessity.

Beer like bread comes from antiquity, and beer comes from bread, because the first beers were made from bread that was thrown in a pot of water and allowed to ferment by the action of naturally occurring air
10 borne yeast. Since barley was the bread of antiquity, and the bread of the people, and beer was the beverage of the people, the loaf of bread that was thrown in the pot was most often made of barley. Right from the beginning (Ancient Egypt) beer was considered a health beverage. Indeed much of the barley consumed by Roman legionnaires and gladiators was in
15 the soluble form of beer made from barley beer loaves made from partially germinated (malted) grain. Use of low alcoholic content stout beer, which contained high levels of viscous solubles as an accepted popular health tonic consumed by the entire family continued in Europe, Africa, Mid East, and America into the first half of the 20th Century. With the
20 development of low viscosity recreational beer in the 20th Century the use of beer as a health tonic has essentially ceased for the general public in urbanized societies. Thus, the beneficial health effects of barley grain were recognized from antiquity. It was also recognized that these benefit were retained when barley was consumed in a soluble form as beer and
25 was associated with the viscosity of the brew, when the viscosity of beers disappeared so did the perceived health benefits. Viscous beta glucan is the active ingredient in healthful barley breads and beers. An especially potent form of viscous beta glucan, long chained beta glucan derived from specially selected waxy hulless and wax hulless short awn barleys has been

described by Fox in U.S. Patent # 5,614,242. The present invention involves high viscosity ingredients (flakes and kibbles), which can be used as cereals and food ingredients to widely extend the use of healthful long chained viscous beta glucan. Special processing techniques are employed to insure that the long chained beta glucans are preserved through processing.

Other patents have described the use of waxy barley for the production of cereal flakes, Alexander (U.S. Patent No. 5,360,619) and Lewis et al. (U.S. Patent No. 5,391,388) incorporated herein. Alexander describes a dry heat stabilized flake, that is used for ready to eat food products and food ingredients. Lewis, et al. describe an uncooked whole flake cereal. Only within this invention is specially selected High Vee™ waxy hulless and waxy hulless short awn barley with high levels of long chained beta glucan resulting in high levels of beta glucan viscosity, processed in such a manner with moist heat to preserve the long chained beta glucan chains, and the associated clinical (hypocholesterolaemic) functionality in a highly digestible fully cooked product.

Summary of The Invention

The present invention is a high viscosity barley flake containing high levels of clinically active long chained, viscous, beta glucan. This product can be used directly as a hot cereal like oatmeal or can be used as an ingredient in a wide array of cooked or baked foods.

The method for producing the high viscosity barley flake (High Vee™ Flake) includes the following steps:

1. Selecting (High Vee™) barley grain having a waxy hulless or waxy hulless short awn genotype, and beta glucan viscosity greater than 15 cps, and long chained beta glucan concentration greater than about 1.0×10^5 / ml of a 5% aqueous meal suspension.

2. Pearling said barley grain in such a manner that about 10% to 20% of the seed including bran and germ layer is abraded away.
3. Conducting vomitoxin test and selecting only pearled grain that has less than 1 ppm vomitoxin.
4. Cutting (kibbling) selected grain into 1 mm 3 mm square bits (kibbles).
5. Steam cooking the kibbles for 15 to 20 minutes at 212 to 240 degrees Fahrenheit.
6. Rolling the cooked kibbles to a thickness ranging from .007 inches to about .0030 inches.
7. Drying the resultant flakes in such a manner that a moisture level below 12% is obtained with drying temperatures never exceeding 50 degrees Centigrade.

Detailed Description of the Invention

In order to provide an understanding of several of the terms used in the specification and claims, the following definitions are provided:

Beta Glucan - the term beta glucan (beta 1-4/1-3 glucosyl pyranose polymer) is intended to refer to the name of a non starchy polysaccharide in which individual glucose molecules (20,000 - 1,000,000) are linked by beta 1-4 and beta 1-3 linkages. Beta Glucan is soluble in warm water (40-45 degrees Centigrade), cellulose is insoluble in water. Beta glucan is the main structural material in the cell walls of barley and oat grain.

Beta Glucan Chain Analysis - As used herein, the term beta glucan chain analysis is intended to refer to a microscopic test in which the actual length and concentration of beta glucan chain fiber is measured with a microscope (100X - 400X).

Beta Glucan Viscosity - the term beta glucan viscosity describes the friction that is created in a solution by the presence of beta glucan chains (fibers) and is measured in cps. All beta glucan viscosities reported herein are measured

on a 5% aqueous suspension of ground grain or ground grain fractions.

5 Centipoise Units (cps) - the term Centipoise Units or (cps) is the name of the units commonly used to measure viscosity. By definition the fundamental unit of viscosity measurement is the "Poise", which is a material requiring a sheer stress of one dyne per square centimeter to produce a sheer of one inverse second, which has a viscosity of one poise or 100 centipoise.

10 Chain Length - the term chain length is a measure of beta glucan fiber length in micron, millimeters, or centimeters. Such measurements can be accomplished by mounting a measured portion (1/10th of a drop) of a beta glucan solution on a calibrated slide and viewing under 100X to 400X magnification. Long beta glucan chains are 1mm-6mm in length and 2-10 microns in diameter.

15 Drying Temperatures - the term drying temperatures refers to the ambient air temperature either naturally or artificially applied to seed as it dries to ripened grain (14% to 7% moisture). Commercial driers reach temperatures exceeding 200 degrees Fahrenheit on circulating grain. The actual seed temperature depends on exposure time and moisture content of the seed when exposed to these drying temperatures.

20 Fully Cooked Cereal or Grain - the term fully cooked cereal or grain is a condition in which more than 90% of the starch granules have pasted rendering them digestible in the human intestine.

High Vee™ Barley - the term High Vee™ Barley is a waxy hulless or waxy hulless short awn barley with beta glucan viscosity greater than about 15 cps and long chained beta glucan concentration greater than about 1.0×10^5 /ml of a 5% aqueous flour suspension.

25 High Viscosity - The term high viscosity is defined as having more than about 15 cps.

Instant Cereal Flake - the term instant cereal flake is a cereal flake, which upon addition of boiling water, and about one minute of cooling, is ready to eat.

30 Long Chained Beta Glucan Fibers - the term long chained beta glucan fibers describes beta glucan fibers that are about 1mm to about 6mm in length and about 2 to about 10 microns in diameter.

Low Viscosity - the term low viscosity is defined as having less than about

5 cps.

5 Pasted (Starch) Granules - the term pasted (starch) granules refers to starch granules that subsequent to heating in a water medium at temperatures between 150 degrees Fahrenheit to 212 Fahrenheit are seen under microscopic evaluation to have cracks, internal fissures, holes, ruptures, and full breaks, which renders the granules susceptible to chemical, enzymatic, and mechanical degradation. It also renders them digestible in the human intestine. Intact granules are largely indigestible.

10 Pearling - the term pearling refers to a form of incomplete milling in which the grain is rotated (tumbled) over an abrasive surface in order to remove a portion of the outer seed layers or hulls. Hulled grain like barley or rice are routinely pearled to remove the adhering hulls. Hulless grains like wheat corn or hulless barley can be pearled to remove the bran layer or the pericarp/aleurone seed layer, prior to grain milling, resulting in a cleaner flour with less bran.

15 Quick Cereal Flake - the term quick cereal flake is a cereal flake, which is added to water and boiled for 1 to 3 minutes, allowed to cool, and then eaten.

Steam cooking - the term steam cooking is the preparation of food suitable for human consumption with wet heat at atmospheric pressure 212 degrees Fahrenheit, or under pressure at boiling points above 212 degrees Fahrenheit.

20 Tempering - the term tempering is the addition of water to grain often above normal storage moisture in order to render the grain more malleable or otherwise suitable for milling, flaking or other forms of grain processing.

25 Viscosity - the term viscosity is the measure of fluid friction. A highly viscous material is one that possesses a great deal of internal friction, it will not pour or spread as easily as material of lesser viscosity.

Whole (grained) Flour - As used herein, the term whole (grained) flour describes a whole meal derived from a grain that has been ground by any number of methods.

30 The High Vee™ barley grain of the present invention, (described in part in U.S. Patent #5,614,242, incorporated herein) can be pearled, kibbled, steam cooked, flaked, and dried to yield cereal flakes in which

there is essentially no damage or breakage to the long chained beta glucan fibers contained in the endosperm cell wall, thus preserving full clinical functionality.

It was unexpected that when specially selected High Vee™ barley grain was processed by the novel methods described herein, that an easy to digest instant cereal and food ingredient was formed in which hard to digest starch granules were pasted (ruptured and converted to a digestible form), while the active clinical component, long chained beta glucan, was left intact. The product of the present invention is unique in that it contains long chained beta glucan which instant oatmeal does not, and because of the pearling step, it is far more stable (can be stored 1 year at room temperature with no rancidity). Other patents to produce flakes from waxy or waxy hulless barley have been described but none has described or achieved a fully cooked flake with long chained beta glucan concentrations greater than 1.0×10^5 /ml of an aqueous flour suspension.

One preferred embodiment of this invention is a pearled, steel cut cooked, High Vee™ flake that is suitable as a quick or instant cereal and as a food ingredient to be used in bake goods as suspended flake, in cooked foods such as sauces where it also serves as a thickener, and in meats like hamburger where it serves as a meat extender, and nutritional supplement. The preferred embodiment is described below:

1. High viscosity waxy hulless or waxy hulless short awn barley with beta glucan viscosity and long chained beta glucan concentration greater than about 1.0×10^5 /ml of a 5% aqueous flour suspension is selected for processing.
2. Pearling said grain in such a manner that 10 to 20% of the seed is abraded away with the more than 50% of the germ and bran (testa and pericarp) removed.
3. Vomitoxin testing conducted on the pearled grain to

insure that vomitoxin levels are less than 1 ppm on a dry weight basis.

4. The pearled vomitoxin screened seed is kibbled (cut) by corrugated steel rolls in roughly square pieces that are 1mm to 3mm in size.
5. The kibbles are steamed or otherwise wet cooked for 15 to 20 minutes until more than 90% of the starch granules have pasted (are cracked or broken, rendering the starch digestible in the human stomach and intestines).
6. The cooked kibbles are flaked with smooth roller mills into a flake that is .007 to .016 inches in thickness.
7. The flakes are dried at a temperature that ranges from 0 to 50 degrees centigrade, with about 26 degrees Centigrade being the optimum.

Another embodiment of this invention is the production of high viscosity (High Vee™) kibbles which can be used in food products in which a grittier mouth feel is desired. One such example is hamburger in which the kibble particle size matches the meat particle size and matches the mouth feel of hamburger. In order to achieve this, the process is halted after completion of the kibbling process (step 4) and the vomitoxin screened kibbles are removed as end product.

Another embodiment of this invention is the production of high viscosity (High Vee™) cooked kibbles which can be used in the same sort of products that are described above in the second embodiment with the added benefit of a more digestible cooked kibble. In order to achieve this, the process is taken through the steam cooking of the kibbles (step 5) and then moved directly to the drying step (step 7).

Since many of the potential users of High Vee™ are middle aged to elderly persons, and many could be sufferers of cancer, heart disease,

arthritis, or other ailments, a highly digestible product is desirable and will aid in the acceptance of High Vee™ flakes by the public. One surprising outcome of all 3 embodiments is that the beta glucan viscosity in the end product (kibbles, cooked kibbles, cooked flakes) increase by approximately 50% to 100% over the starting materials due in part to the fact that the pearlings remove much insoluble fiber contained in the pericarp and testa which adds nothing to the soluble beta glucan content, thus increasing the soluble long chained beta glucan content by simple dilution effect. It is also possible that the inner layers of the seed (endosperm) contain a higher concentration of long chained viscous beta glucan. It should be noted that there is no tempering step (addition of water to grain or pearled grain) in the described embodiments. This procedure is often employed prior to flaking or kibbling but hydration has the effect of reducing the length of long chained beta glucan. Unexpectedly it has been found that when using high viscosity long chained waxy hulless or waxy hulless short awn barleys at storage moisture (6% to 13.5%), that steam cooking is a sufficient preparation for flaking or kibbling.

Also unexpectedly, it was found that by steam cooking untempered bits of grain (kibbles) a fully cooked (digestible), flake was produced in which the long chained beta glucans were preserved, with the degrading cereal beta glucanases (endogenous enzymes that degrade beta glucan) inactivated. It should be noted that neither the Alexander nor Lewis et al. processes have a kibble step before flaking, and both use tempered grain as starting material; both steps are key components to the process described herein.

It was determined that select High Vee™ waxy hulless or waxy hulless short awn barley, with a clean bright appearance and vomitoxin levels no greater than 1ppm can be used in the above described process/invention, as a whole grain, with no health problems (associated

with vomitoxin), and with a one year shelf life as described in the preferred embodiment with the result that a whole grain High Vee™ cooked flake, and kibble, (cooked and uncooked) can be produced. As with the pearled High Vee™ starting material long chain beta glucan
5 contained in the whole grain waxy hulless and waxy hulless short awn barley grain is preserved throughout the steam cooking process.

EXAMPLES

The method of this invention is illustrated below in the following examples. These examples are submitted for illustrative purposes only and
10 are meant in no way to limit the scope of the invention.

EXAMPLE 1

A sample of high beta glucan viscosity waxy hulless barley with a beta glucan viscosity of 20 cps, a concentration of long chained beta glucan that is greater than 1.0×10^5 /ml of a 5% aqueous flour suspension,
15 and a vomitoxin concentration greater than 4 ppm is selected for processing. The barley grain is pearled so that about 15% of the seed weight is abraded away. The pearled grain is sieved over a 5/64 inch round holed sieve with the overs retained and vomitoxin ppm reduced to an acceptable level of less than 1 ppm. The resultant pearled grain is steel
20 cut in manner well known to skilled artisans to produce kibbles that are in the form of cubes that are 1 mm to 2 mm on a side. The beta glucan viscosity increased to 40 cps, with increased long chained beta glucan concentration in these kibbles. A sample of the above described kibble was blended with hamburger, 15 parts kibble to 85 parts hamburger, and
25 prepared and cooked in the traditional manner as patties, and found to yield an extremely juicy and tasty patty with reduced animal fat content.

EXAMPLE 2

A sample of high beta glucan viscosity, waxy hulless barley with a beta glucan viscosity of 33 cps, a concentration of long chained beta

glucan that is greater than 1.0×10^5 /ml of a 5% aqueous flour suspension, and a vomitoxin concentration greater than 6 ppm. The barley grain is pearled so that about 20% of the seed weight is abraded away. The pearled grain is sieved over a 5/64 inch round holed sieve and the overs retained and the vomitoxin ppm reduced to an acceptable level of less than 1 ppm. The resultant pearled grain is steel cut using corrugated rollers in a manner that is well known to those skilled in the art to produce kibbles that are in the form of cubes that are 1 mm to 2 mm on a side. The kibbles are steamed cooked for 15 minutes and dried at 26 degrees to a moisture concentration of 12.0%. The beta glucan viscosity of the resultant kibble was 50 cps and long chained beta glucan concentration increased. When incorporated into hamburg at concentrations from 5% to 20% the kibbled burger was found to yield an extremely juicy and tasty patty with reduced animal fat content.

15

EXAMPLE 3

A sample of high beta glucan viscosity waxy hulless barley with a beta glucan viscosity of 17 cps, a concentration of long chained greater than 1.0×10^5 /ml of a 5% aqueous flour suspension, and a vomitoxin concentration greater than 3 ppm. The barley is pearled so that about 15% of the seed weight is abraded away, said pearled grain is sieved over a 5/64 inch round holed sieve with the overs retained and vomitoxin ppm reduced to an acceptable level of less than 1 ppm. The resultant pearled grain is steel cut in a manner well known to those skilled in the art to produce kibbles that are in the form of cubes 1 mm to 2 mm on a side. The kibbles are steamed cooked for 20 minutes at 210 to 240 degrees Fahrenheit with the temperature ascending with time until more than 90% of the starch granules within the kibbles have pasted. The cooked kibbles are immediately flaked with smooth roller mills in a manner well known in the art to form flakes that are .007 to .016 inches thickness. The flakes

are then immediately dried at about 26 degrees Centigrade with constant aeration. Beta glucan viscosity of the flakes resultant increases to 30 cps, and the concentration of long chained beta glucan is maintained. The resultant high viscosity flakes can be used as an instant or quick cereal

5 flake. The distinction between instant and quick flakes is strictly based on particle size (thickness). Instant flakes are in the .007 to .010 inch range and quick flakes are in the .011 to .018 inch range. Since the flakes have already been cooked as kibbles the distinction between instant and quick flakes is based solely on speed of hydration, with smaller particles (instant

10 flakes) hydrating more rapidly than larger (quick flakes) particles. Often it is necessary to produce quick and instant flakes as separate smooth roller mill runs, setting the rollers tighter for the thinner flakes. However, in this example a range of thicknesses were produced between .007 inches to .016 inches and the two flake types were produced (separated) by sieving.

15 The instant/quick flakes can be used not only as a hot cereal, but also as an ingredient in baked goods such as breads, cookies, and muffins. The instant/quick flakes can also be blended with meat (meat loaf), can be used in sauces as a healthful thickener (gravies, tomato sauces), and can be used in any soup as an healthful thickener.

20

EXAMPLE 4

A sample of high beta glucan viscosity waxy hulless barley with a beta glucan viscosity of 30 cps, a concentration of long chained beta glucan greater than 1.0×10^5 /ml of a 5% aqueous flour suspension, and a vomitoxin concentration no greater than 1 ppm were selected. The whole

25 grain kernel is steel cut in a manner well known to those skilled in the art to product kibbles that are in the form of cubes 1mm to 2 mm on a side. The kibbles are steam cooked for 20 minutes at 210 to 240 degrees Fahrenheit with the temperature ascending with time. The cooked kibbles are immediately flaked with smooth roller mills in a manner well known in

the art to form flakes that are .007 to .016 in thickness. The flakes are then dried at about 26 degrees Centigrade with constant aeration, and sieved to yield instant and quick flakes. Vomitoxin testing of the flakes indicated levels of less than 1 ppm, and long chained beta glucan analysis indicated a concentration greater than 1.0×10^5 /ml of an aqueous flour suspension. The instant/quick whole grain flakes can be used not only as a hot cereal but also as an ingredient in baked goods, such as breads, cookies and muffins. The instant/quick whole grain flakes can also be blended with meat (meat loaf) or used in sauces (gravies, tomato sauces) and soups as a healthful thickener.

EXAMPLE 5

A sample of quick style (High Vee TM) whole grain flakes described in Example 4 was added as an ingredient to preparation of Lipton Chicken Noodle soup. This compared to a sample of whole waxy hulless barley flakes which were flaked with just enough steam cooking (plasticizing) to form a flake of about .022 inches in thickness. These whole flakes were uncooked with less than 30% of the starch granules pasted. An additional comparison was made with a commercial sample of quick oat flakes.

Waxy hulless barley flakes contain no amylose (the soluble form of starch in normal starch granules). Thus, any viscosity created as these flakes are added to hot soup is due to the soluble β -glucans. Oats, the other cereal with β -glucan, does not contain waxy starch and upon heating will release both amylose and β -glucan into solution. Thus, viscosity created by the oat flakes can be due to a combination of starch and β -glucan.

The viscosity created by these cereal flakes is an indication of clinical functionality. The higher the quantity and quality of the cholesterol and fat absorbing β -glucan the more viscous the resultant

fortified soup will be. Viscosity created by amylose starch is degraded quickly in the small intestine by starch degrading enzymes naturally present in the saliva and has no clinical functionality.

The methods and results of this experiment are listed below. The High Vee™ Flakes are superior to the whole, uncooked barley flakes at every concentration level. They are also clearly superior to the quick oat flakes at the ¾ oz addition level. Much if not all the viscosity of the oat flakes was due to the milky colored amylose that was observed in the soup. The oats created a 'milky' mess. The whole, uncooked barley flakes were apparent as large, hard-to-digest objects in the soup. The High Vee™ flakes at the ¼ to ½ addition level were barely detectable and easily consumed and digested. The High Vee™ flakes created a smooth, appealing mouthfeel and no aftertaste - they were basically bland and undetectable to most who sampled the soup. In addition, there was no characteristic linseed smell or taste associated with grain rancidity which is remarkable since these High Vee™ whole grain flakes were 1 year old with no special storage conditions (room temperature, 26°C).

Experiment 1

Purpose: To compare High Vee™ flakes with whole barley flakes (WBF) and quick oat flakes.

Methods Used:

Viscosity in soup

- Boiled Lipton Soup mix with 4 C water; simmered for 5 minutes
- Weighed 1 oz (28.4g), ¾ oz (21.3g), ½ oz (14.2g) and ¼ oz (7.1g) of each sample into individual canning jars;
- Poured 1 C of soup (not noodles) into each jar (3 jars/batch of soup) and covered;
- Allowed to sit for 40 min, then cooled to 26°C and read the viscosity**

**Viscosity measurement - liquid was poured into a 25 ml graduated cylinder and 10 cm marked. Time (in seconds) for ball to travel 10 cm is recorded. Each second of travel time is equivalent to a cps viscosity unit.

5

Amount	Liquid/solid	Viscosity (sec)		
		High Vee TM	WBF	Oats
1 oz	8.3	240	6	-
³ / ₄ oz	11.1	20	3	10
¹ / ₂ oz	16.7	15	3	-
¹ / ₄ oz	33.3	3	2	-
water	-	1	1	-

10

This is just one example of how High Vee TM flakes are used as a versatile food ingredient to improve the nutritional functionality of food products while maintaining good taste and textural properties resulting in products that are both good tasting and better nutritionally.

15

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity and understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the invention, as limited only by the scope of the appended claims

CLAIMS

WHAT IS CLAIMED:

1. A method of producing high viscosity waxy hulless barley flakes with beta glucan viscosity greater than 15 cps and a long chained beta
5 glucan concentration greater than about 1.0×10^5 /ml of a 5% aqueous flour suspension comprising:
 - (a) selecting high beta glucan viscosity barley grain;
 - (b) pearling said high viscosity barley grain;
 - (c) kibbling said pearled high viscosity barley grain;
 - 10 (d) steam cooking the said pearled high viscosity barley kibble;
 - (e) flaking the said pearled high viscosity cooked barley kibble; and
 - (f) drying the said pearled high viscosity cooked barley
15 flake at a temperature below 50 degrees Centigrade.
2. High viscosity waxy hulless barley flakes with beta glucan viscosity greater than 15 cps and a long chained beta glucan concentration greater than about 1.0×10^5 /ml of a 5% aqueous flour suspension, produced by the method of claim 1.
- 20 3. The high viscosity barley flakes of claim 2 wherein the flakes are made from barley having a waxy hulless short awn genotype.
4. A method of producing high viscosity waxy hulless barley kibbles with beta glucan viscosity greater than 15 cps and a long chained beta glucan concentration greater than about 1.0×10^5 /ml of a 5% aqueous
25 flour suspension, the method comprising the following steps:
 - (a) selecting high beta glucan viscosity barley grain;
 - (b) pearling said high viscosity barley grain; and
 - (c) kibbling said pearled high viscosity barley grain;

5. High viscosity waxy hulless barley kibbles with beta glucan viscosity greater than 15 cps and a long chained beta glucan concentration greater than about 1.0×10^5 /ml of a 5% aqueous flour suspension, produced by the method of claim 4.

5 6. The high viscosity barley kibble of a claim 5 wherein the kibble is made from barley having a waxy hulless short awn genotype.

7. A method of producing steam cooked high viscosity waxy hulless barley kibbles with beta glucan viscosity greater than 15 cps and a long chained beta glucan concentration greater than about 1.0×10^5 /ml of
10 5% aqueous flour solution, produced by a process which comprises:

(a) selecting high beta glucan viscosity barley grain;

(b) pearling said high viscosity barley grain;

(c) kibbling said pearled high viscosity barley grain;

(d) steam cooking said pearled high viscosity barley
15 kibble; and

(e) drying said pearled high viscosity cooked barley kibble at a temperature below 50 degrees Centigrade.

8. Steam cooked high viscosity waxy hulless barley kibbles with beta glucan viscosity greater than 15 cps and a long chained beta glucan
20 concentration greater than about 1.0×10^5 /ml of a 5% aqueous flour suspension, produced by the method of claim 7.

9. Steam cooked high viscosity waxy hulless barley kibble of claim 8 wherein the kibble flake is made from barley having a waxy hulless short awn genotype.

25

10. A method of producing whole grain high viscosity waxy hulless barley flakes with a beta glucan viscosity greater than 15 cps and a long chained beta glucan concentration greater than about 1.0×10^5 /ml of a 5% aqueous flour suspension; comprising the steps of:

- (a) selecting high beta glucan viscosity barley grain;
- (b) kibbling said high viscosity barley grain;
- (c) steam cooking the said high viscosity barley kibble;
- (d) flaking the said high viscosity cooked barley kibble;
- 5 and
- (e) drying said high viscosity cooked barley flake at a temperature below 50 degrees Centigrade.

11. Whole grain high viscosity waxy hulless barley flakes with beta glucan viscosity greater than 15 cps and a long chained beta glucan
10 concentration greater than 1.0×10^5 /ml of a 5% aqueous flour suspension, produced by the method of claim 10.

12. The whole grain high viscosity barley flakes of claim 11 wherein the flakes are made from barley having a waxy hulless short awn genotype.

15 13. A method of producing whole grain high viscosity waxy hulless barley kibbles with beta glucan viscosity greater than 15 cps and a long chained beta glucan concentration greater than about 1.0×10^5 /ml of a 5% aqueous flour suspension, the method comprising the following steps:

- 20
- (a) selecting high beta glucan viscosity barley grain; and
 - (b) kibbling said high viscosity barley grain.

14. Whole grain high viscosity waxy hulless barley kibbles with beta glucan viscosity greater than 15 cps and a long chained beta glucan concentration greater than about 1.0×10^5 /ml of a 5% aqueous flour
25 suspension, produced by the method of claim 13.

15. The whole grain high viscosity barley kibble of claim 14, wherein the kibble is made from barley having a waxy hulless short awn genotype.

16. A method of producing steam cooked whole grain high viscosity waxy hulless barley kibbles with a beta glucan viscosity greater than 15 cps and a long chained beta glucan concentration greater than about 1.0×10^5 /ml of a 5% aqueous flour suspension, comprising:

- 5 (a) selecting high beta glucan viscosity barley grain;
 (b) kibbling said high viscosity barley grain;
 (c) steam cooking said high viscosity barley kibble; and
 (d) drying said steam cooked high viscosity barley kibble
 at a temperature below 50 degrees Centigrade.

10 17. Steam cooked whole grain high viscosity waxy hulless barley kibbles with beta glucan viscosity greater than 15 cps and a long chained beta glucan concentration greater than about 1.0×10^5 /ml of a 5% aqueous flour suspension, produced by the method of claim 16.

15 18. Steam cooked whole grain high viscosity waxy hulless barley kibble of claim 17, wherein the flake is made from barley having a waxy hulless short awn genotype.

INTERNATIONAL SEARCH REPORT

Inter. Appl. No.

PCT/US 98/17053

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A23L1/10 A23L1/168 A23L1/308 A61K35/78

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A23L A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 614 242 A (FOX GREGORY J) 25 March 1997 cited in the application see column 3, line 10 - line 65 see column 7, line 44 - line 58 see column 9, line 18 - line 37 see column 10, line 5 - line 36 see examples 6, 11, 12, 15 ---	2-6, 8, 9, 11-15, 17, 18
A	EP 0 606 080 A (QUAKER OATS CO) 13 July 1994 see column 2, line 54 - column 7, line 29 see column 8, line 8 - line 52; example 1 --- -/--	1-18

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

9 November 1998

Date of mailing of the international search report

16/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Vuillamy, V

INTERNATIONAL SEARCH REPORT

Inter. .ional Application No

PCT/US 98/17053

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 393 870 A (BYRON AGRICULTURAL CO) 24 October 1990 see page 2, line 42 - page 3, line 14 see page 4, line 4 - line 10 see examples I-III & US 5 391 388 A cited in the application ---	1-18
A	US 5 360 619 A (ALEXANDER DONALD J) 1 November 1994 cited in the application see column 4, line 7 - column 7, line 26 -----	1-18

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/17053

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5614242 A	25-03-1997	US 5725901 A	10-03-1998
EP 0606080 A	13-07-1994	CA 2112835 A	07-07-1994
		FI 940038 A	07-07-1994
		NO 940036 A	07-07-1994
EP 0393870 A	24-10-1990	AU 624230 B	04-06-1992
		AU 5237790 A	11-10-1990
		CA 2013707 A,C	05-10-1990
		DE 69002368 T	13-01-1994
		US 5391388 A	21-02-1995
US 5360619 A	01-11-1994	AU 7961994 A	01-05-1995
		WO 9509540 A	13-04-1995